

Preventive Medicine



Melatonin Sublingual

DESCRIPTION

Melatonin Sublingual tablets contain 3 mg of purest pharmaceutical grade melatonin, in a convenient sublingual tablet.

FUNCTIONS

Melatonin is a natural hormone nutrient that is synthesized from the amino acid tryptophan by the pineal gland in the back of the brain. Melatonin also occurs in small amounts in a variety of foods. In the body, melatonin appears to regulate sleep/wake cycles, support normal immune function, and protect cells from free radical damage. Most of the research on melatonin has focused on its role in maintaining normal sleep/wake rhythms. The perception of daylight in the eyes is a signal for the pineal gland to inhibit melatonin synthesis and release. At night or in the dark, the body's melatonin production rises. The rise in plasma melatonin is thought to be responsible for bringing on sleep. Nocturnal melatonin production is highest in children and begins to decline from adolescence on until it is virtually absent in the elderly. Melatonin supports normal immune function by helping maintain the activity of circulating natural killer cells. It also has been found to function as an antagonist for stress-induced immunosuppression. Melatonin is considered a potent antioxidant that enters all body cells to help prevent free radical damage. In the brain, melatonin is perhaps the most important physiological antioxidant. Due to its lipid and water-soluble properties, it can freely cross the blood-brain barrier. In vitro studies show that melatonin is more effective than glutathione in scavenging toxic hydroxyl radicals, and also more efficient than vitamin E in neutralizing peroxy radicals which can induce DNA damage. Furthermore, melatonin stimulates the main antioxidant enzyme of the brain, glutathione peroxidase.

INDICATIONS

Melatonin Sublingual tablets may be a useful dietary supplement for individuals wishing to supplement their diet with melatonin.

FORMULA (#81322)

Each Sublingual Tablet Contains:
Melatonin 3 mg

SUGGESTED USE

Adults take 1 tablet daily or as directed by physician.

SIDE EFFECTS

Do not use before or while operating a motor vehicle or heavy machinery.

STORAGE

Store in a cool, dry place, away from direct light. Keep out of reach of children.

DISTRIBUTED BY

Preventive Medicine
Carol Joy Gardner, D.O.
905 Roosevelt Hwy, #210, Colchester Vt 05446
802-879-6544

REFERENCES

Alvarez B, Dahlitz M, Vignau J, Parkes JD: The delayed sleep phase syndrome: Clinical and investigative findings in 14 patients. *J Neurol Neurosurg Psychiatry* 1992;55:665-670.
Armstrong SM, Redman JR: Melatonin: A chronobiotic with anti-aging properties? *Med Hypotheses* 1991;34:300-309.
Beck-Friis J, Kjellman BF, Aperia B et al: Serum melatonin in relation to clinical variables in patients with major depressive disorder and a hypothesis of a low melatonin syndrome. *Acta Psychiatr Scand* 1985;71:319-330.
Bismar K, Morgensen L, Wetterberg L: Depressed melatonin secretion in patients with nightmares due to betaadrenoceptor blocking drug. *Acta Med Scand* 1987;221:155-158.
Dahlitz M, Alvarez B, Vignau J et al: Delayed sleep phase syndrome response to melatonin. *Lancet* 1991;337:1121-1124.
Lieberman HR: Behavior, sleep and melatonin. *J Neural Transm Suppl* 1989;21:233-241.
Lissoni P, Barni S, Cattaneo G et al: Clinical results with the pineal hormone melatonin in advanced cancer resistant to standard antitumor therapies. *Oncology* 1991;48:448-450.

**These supplements have not been evaluated by the Food and Drug Administration.
This product is not intended to diagnose, treat, cure or prevent any disease.**