

Preventive Medicine



Saccharomyces Boulardii

DESCRIPTION

Saccharomyces boulardii is a non-pathogenic, transient yeast, long-used for diarrhea or dysentery.* It is a hardy, acid-resistant, temperature tolerant microorganism that is not affected by anti-bacterial agents.*

FUNCTION

S. boulardii is genetically and functionally distinct from brewer's yeast (*S. cerevisiae*) and different from pathogenic *Candida* species. *S. boulardii* increases the activities of intestinal brush border enzymes such as disaccharidases, a-glucosidases, alkaline phosphatases, and aminopeptidases. It secretes a leucine aminopeptidase that appears to support against allergies to dietary proteins following acute gastroenteritis.* *S. boulardii* augments the intestinal absorption of D-glucose coupled with sodium that may enhance uptake of water and electrolytes during diarrhea.* It increases stool concentrations of short-chain fatty acids that nourish colon mucosal cells.* It modulates immunity by boosting intestinal secretion of IgA and increasing crypt cell immuno-globulin receptors.* It inhibits inflammatory cytokine pathways and secretes a factor that blocks *C. difficile* toxin A activities.* Research has found that *S. boulardii* supports healthy intestinal function in an array of clinical settings including antibiotic-associated diarrhea, *C. difficile* disease, inflammatory bowel disease, irritable bowel syndrome, and traveler's diarrhea.* Some of the key functions of *S. boulardii* include:

- Support the normal gastrointestinal flora during antibiotic therapy.*
- Assists individuals with diarrhea, including antibiotic-associated diarrhea.*
- Enhances the number of healthful bifidobacteria in the colon while

simultaneously suppressing populations of pathogenic clostridia.*

- Mitigates the effects of toxins secreted by a number of common intestinal pathogens.*
- Produces antifungal toxins and organic acids known to naturally inhibit or kill disease-causing *Candida* yeast species provides support against *C. difficile*, toxigenic *E. coli*, and other gastrointestinal pathogens.*
- Helps stimulate host immune defenses and improves gut barrier function.*

INDICATION

S. boulardii is generally used for short-term gastrointestinal support. A common supplement routine is three to four weeks or up to two weeks after diarrhea has resolved.

FORMULA

Serving Size: 1 capsule

Saccharomyces boulardii 150 mg

*This product is free of artificial flavors, preservatives, colorings and the following common allergens: milk, eggs, fish, shellfish, tree nuts, peanuts, wheat/gluten, corn and soybeans. People with known allergies to *Saccharomyces cerevisiae* and individuals with chronic indwelling catheters should avoid use of *S. boulardii*.

SUGGESTED USE

As a dietary supplement take one capsule twice daily with meals or as directed by your health care provider.

STORAGE

Store in the refrigerator once opened.

DISTRIBUTED BY

Preventive Medicine
Carol Joy Gardner, D.O.
905 Roosevelt Hwy, #210, Colchester, Vt 05446
802-879-6544

*These statements have not been evaluated by the Food and Drug Administration.
This supplement is not intended to diagnose, treat, cure or prevent any disease.

Formula # V776-06